

Public ref
gathering
this collec
Davis Hig

AD-A228 957

hour per response, including the time for searching existing data sources, collection of information. Send comments regarding this estimate or any other aspect of Shington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson
ment and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. Agency

1990

3. Report Type and Dates Covered.

Abstract

4. Title and Subtitle.

A 1/8 Degree Model of the North Pacific Ocean

5. Funding Numbers.

Program Element No. 61153N

Project No. 3103

Task No. 030

Accession No. DN258028

6. Author(s).

H. E. Hurlburt, E. J. Metzger, A. J. Wallcraft,
and P. J. Hogan

7. Performing Organization Name(s) and Address(es).

Naval Oceanographic and Atmospheric
Research Laboratory
Stennis Space Center, MS 39529-50048. Performing Organization
Report Number.

AB 90:323:113

9. Sponsoring/Monitoring Agency Name(s) and Address(es).

Naval Oceanographic and Atmospheric
Research Laboratory
Stennis Space Center, MS 39529-500410. Sponsoring/Monitoring Agency
Report Number.

AB 90:323:113

11. Supplementary Notes.

*Continued on next page
EOS

12a. Distribution/Availability Statement.

Approved for public release; distribution
is unlimited.

12b. Distribution Code.

13. Abstract (Maximum 200 words).

Results from a 1/8 degree Pacific Ocean model (.125 degree, .176 degree resolution lat, long for each variable are compared with results from several versions of a model with 1/4 degree resolution. All of the models are layered primitive equation, cover the Pacific Ocean north of 20S, include marginal seas, and have a free surface. The 1/8 degree model has 2 layers and realistic topography. The versions of the 1/4 degree model are 1.5 and 3.5 layer reduced gravity and 2-layer finite depth with realistic topography or a flat bottom. These models are used to show (1) the effects of horizontal resolution and vertical structure on flow instabilities and (2) the relative importance of time-varying wind and flow instabilities in producing a time dependent oceanic response. For example, flow instabilities are most widespread in the 3.5 layer reduced gravity model. The barotropic mode has the largest impact on flow instabilities in the Kuroshio region. A salient difference between the 1/8 degree model and the corresponding 1/4 degree model the maximum occurs near the end of the inertial jet between 155 degree E and 160 degree E. In the 1/8 degree*

14. Subject Terms.

(U) Ocean Models; (U) Ocean Forecasting;
(U) Fronts (Oceanography)

15. Number of Pages.

16. Price Code.

17. Security Classification
of Report.

Unclassified

18. Security Classification
of This Page.

Unclassified

19. Security Classification
of Abstract.

Unclassified

20. Limitation of Abstract.

SAR

*model they are much more widespread, extending from south of Japan to 160 degree E and northward to about 40 degree N and the subarctic front, more like the RMS variability patterns from GEOSAT altimetry. All of the models simulate the basic features of the North Pacific, including the subtropical and subpolar gyres, the major current systems (Kuroshio, Oyashio, North Equatorial, Mindanao, North Equatorial Countercurrent) and associated fronts including the subarctic and the Kuroshio extension. Only the models with realistic topography exhibit the bifurcation of the Kuroshio at the Shatsky rise and only the 3.5-layer reduced gravity model simulates the equatorial undercurrent.

INSPECTED

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

031F-5 0945h

A 1/8° Model of the North Pacific Ocean

H E Hurlburt and E J Metzger (Both at: Naval Oceanographic and Atmospheric Research Laboratory, Code 323, Stennis Space Center, MS 39529-5004; 601-688-4626; GTE Mail H.Hurlburt)
A J Wallcraft (JAYCOR, Bldg 1103, Stennis Space Center, MS 39529)
P J Hogan (Sverdrup Technologies, Bldg 2109, Stennis Space Center, MS 39529)

Results from a 1/8° Pacific Ocean model (.125°, .176° resolution lat, long for each variable) are compared with results from several versions of a model with 1/4° resolution. All of the models are layered primitive equation, cover the Pacific Ocean north of 20S, include marginal seas, and have a free surface. The 1/8° model has 2 layers and realistic topography. The versions of the 1/4° model are 1.5 and 3.5 layer reduced gravity and 2-layer finite depth with realistic topography or a flat bottom.

These models are used to show (1) the effects of horizontal resolution and vertical structure on flow instabilities and (2) the relative importance of time-varying winds and flow instabilities in producing a time dependent oceanic response. For example, flow instabilities are most widespread in the 3.5 layer reduced gravity model. The barotropic mode has the largest impact on flow instabilities in the Kuroshio region. A salient difference between the 1/8° model and the corresponding 1/4° model is the distribution of the eddy generation and oceanic variability in the Kuroshio region. In the 1/4° model the maximum occurs near the end of the inertial jet between 155°E and 160°E. In the 1/8° model they are much more widespread, extending from south of Japan to 160°E and northward to about 40°N and the subarctic front, more like the RMS variability patterns from GEOSAT altimetry. All of the models simulate the basic features of the North Pacific, including the subtropical and subpolar gyres, the major current systems (Kuroshio, Oyashio, North Equatorial, Mindanao, North Equatorial Countercurrent) and associated fronts including the subarctic and the Kuroshio extension. Only the models with realistic topography exhibit the bifurcation of the Kuroshio at the Shatsky rise and only the 3.5-layer reduced gravity model simulates the equatorial undercurrent.